



## Efficiency of Nano-chitosan and Azotobacter on growth and yield of kohlrabi plant

Hayder. A. A. Turk<sup>1</sup> Mahmood.T.Mahmood<sup>2</sup> Mazin.M. A. Ameen<sup>3</sup> Radhiyah. A. H. Ahmad<sup>4</sup> Abdel Rahman .M. Al Tawaha<sup>5</sup>

<sup>1</sup>Department of Science. Open Educational College , Iraq.

<sup>2</sup>Karbala Agriculture Directorate, Ministry of Agriculture, Iraq.

<sup>3</sup>Shiit Endowment Diwan /Imam Al-Kadhum College, Iraq.

<sup>4</sup>College of Education for Pure Sciences. Ibn Al-Haitham, University of Baghdad, Iraq.

<sup>5</sup>Department of Biological Sciences, Al-Hussein Bin Talal University, Amman, Jordan.

\*Correspondence: [haidrturk1967@gmail.com](mailto:haidrturk1967@gmail.com)

### KEY WORDS:

Bio fertilizer, Azotobacter, Nano-Chitosan, Kohlrabi

Received: 17/01/2025

Revision: 22/03/2025

Proofreading: 14/05/2025

Accepted: 28/05/2025

Available online: 30/09/2025

© 2025. This is an open access article under the CC by licenses <http://creativecommons.org/licenses/by/4.0>



### ABSTRACT

The research was conducted in one of the fields of the Department of Plant Production in the desert region (tomato development project) affiliated with the Directorate of Agriculture of the Holy Karbala Governorate for the purpose of studying the effect of spraying the shoots of the Kohlrabi plant (hybrid) with the Nano biopolymer (Chitosan) and the *Azotobacter* bio inoculum on the roots of the plant seedlings in terms of growth indicators and yield for the autumn agricultural season. 2023-2024 ,The means were compared according to the Duncan multiple ranges test at significant to level of 0.05.with R.C.B.D . The first factor was spraying the shoots with Nano-Chitosan at a concentration of (0, 1, and 2) g L<sup>-1</sup>, two weeks after transplanting, and two sprays every 14 days. The second factor has three levels, the first is planting without a vaccine, the second is inoculating the seedling roots with *Azotobacter* at a level of 5 g , and the third level is inoculating with 10 grams of the biological factor. The results showed a significant superiority of the interaction treatment between the bio-addition of *Azotobacter* at a level of 5 g .and a concentration of 2 g of Nano-chitosan in the characteristics of each of the plant length 60 cm, the number of leaves 21.67 leaves. plant<sup>-1</sup>the leaf content of chlorophyll 77.59 mg 100 g<sup>-1</sup>, The Weight of the knobs cm plant<sup>-1</sup> 278.9 g , the total yield 14.87 ton . h<sup>-1</sup>, content of nitrogen 3.43% and the knobs content of protein 21.48% compared to the control treatment, which gave the lowest value.

## كفاءة الكيتوسان النانوي و الازوتوباكتري في نمو وإنتاجية نبات الكل

حيدر عبد المنعم عبد الامير ترك<sup>1</sup> ، محمود طالب محمود<sup>2</sup> ، مازن موسى عبد امين<sup>3</sup> ، رضية علي حسن احمد<sup>4</sup> ، عبد الرحمن محمد الطحاوي<sup>5</sup>

قسم العلوم , الكلية التربوية المفتوحة , العراق  
مديرية زراعة محافظة كربلاء المقدسة , العراق  
ديوان الوقف الشيعي , كلية الامام الكاظم , العراق  
كلية التربية للعلوم الصرفة – بن الهيثم , جامعة بغداد , العراق  
قسم علوم الحياة , جامعة الحسين بن طلال , الاردن

### الخلاصة

أجري البحث في أحد حقول قسم الإنتاج النباتي في المنطقة الصحراوية (مشروع تنمية الطماعة) التابع لمديرية زراعة محافظة كربلاء المقدسة لغرض دراسة تأثير رش نبات الكلم (الهجين) بالبوليمر الحيوي النانوي (الكيتوسان) والملح الحيوي الازوتوباكتري على جذور شتلات النبات في مؤشرات النمو والحاصل للموسم الزراعي الخريفي 2023-2024، وقورنت المتوسطات وفق اختبار دنكن متعدد المدى عند مستوى معنوي 0.05 ، وكان العامل الأول هو رش النباتات بالنانو كيتوسان بتركيز (0، 1، 2) غم لتر<sup>-1</sup>، بعد أسبوعين من الزراعة، ورشتين كل 14 يومًا. العامل الثاني له ثلاثة مستويات، الأول الزراعة بدون لقاح، والثاني تلقيح جذور الشتلات الازوتوباكتري بتركيز 5 غم، والثالث تلقيحها بـ 10 غم من العامل الحيوي. وقد أظهرت النتائج تفوقًا معنويًا لمعاملة التفاعل بين الإضافة الحيوية للازوتوباكتري بتركيز 5 غم وتركيز 2 غم من النانو كيتوسان في صفات كل من طول النبات 60 سم، وعدد الأوراق 21.67 ورقة نبات<sup>-1</sup>، ومحتوى الأوراق من الكلوروفيل 77.59 ملغم/100 غم<sup>-1</sup>، ووزن الكورمات غم نبات<sup>-1</sup> 278.9 غم، والمحصول الكلي 14.87 طن/هـ<sup>-1</sup>، ومحتوى النيتروجين 3.43%، ومحتوى الكورمات من البروتين 21.48%، وذلك مقارنةً بمعاملة المقارنة التي أعطت أقل قيمة.

**الكلمات المفتاحية:** السماد الحيوي ، ازوتوباكتري ، الكيتوسان النانوي ، الكلم .

### INTRODUCTION

Plants of the vegetable crop family gain the utmost importance in the human diet because of the nutritional and health value they represent, including the Kohlrabi *Brassica oleracea var gongylodes* , which belongs to the *Brassicaceae* family, as the edible part of the plant is the swollen stem, and the plant contains mineral elements, vitamins, effective compounds beneficial to public health, and antioxidants. the plant has several varieties, including local ones that are elongated and other types that are balled. Its uses are in cooking, salads, or making pickles (Ali and Khalid, 2023).

The steady increase in population growth in the world has led to an increase in demand for food products to meet the actual need (Al-Mosawy, 2023), which has led agricultural producers to excessive use of chemical fertilizers in addition to the use of intensive agriculture by exploiting the entire land area, which has led to the deterioration of the soil's nutritional stock by a rate approaching between (5-50) %. The increase in levels of chemical fertilizers has led to the accumulation of harmful compounds in the human body as a result of consumption and the emergence of health problems as a result of that accumulation (Turk et al, 2020). Also, the use of bio fertilizers, which depends on the use of natural biological systems to transfer important nutrients to the plant without resorting to harmful chemical fertilizers, with the aim of maintaining the level of production so that the food becomes healthy and safe (Mohammed et al, 2017).

The need for the emergence of biological alternatives, including bio fertilizers, aims to reduce the use of traditional fertilizers. Organic farming reduces environmental pollution

problems and works to sustain agriculture by increasing metabolic activities to revive soil microbiology (Ahmed, 2016), which contributes to providing the plant with elements such as phosphorus, which also plays a major role in the synthesis of energy-rich compounds that the plant acquires to build carbohydrates and enzymes (Taiz and Zeiger, 2014). Microorganisms provide a suitable environment for activating the vital functions of plants, which increases vegetative growth. These organisms are more beneficial to the soil than modern fertilizers, as they contribute to increasing the soil's mineral and ion content and improving the chemical condition, in addition to the metabolic activities they perform. (Al-Sharia, 2022). They also contribute to increasing the concentration of nitrogen in the soil content and thus accelerating plant growth. *Azotobacter* bacteria contribute to improving the root environment and work to provide the root zone with important nutrients for the purpose of increasing the plant's absorption capacity through the readiness of these ions in the soil solution (Al-Bayati *et al*, 2023).

Clean agriculture works to manage natural biological systems in the transfer of important nutrients to plants without resorting to harmful chemical fertilizers, with the aim of maintaining the level of production and keeping plants free of pollutants so that food becomes healthy and safe (Sabalpara and Lalit, 2016). It's also achieves increased production and environmental protection to produce healthy and safety food (Turk., 2023).

The *Azotobacter chroococcum* bacteria It is characterized by its ability to fix nitrogen, in addition to acting as growth promoters, as it has a role in increasing the growth and productivity of plants (Turan *et al.*, 2006). Bio fertilizers are used to reduce the addition of chemical fertilizers by no less than 25%, in addition to their role in reducing Environmental pollution problems and works to sustain agriculture (Al-Juthery *et.al*, 2020). Diversifying the sources of organic fertilizers has prompted specialists in the agricultural fields to search for alternatives that can be used in organic agriculture, and among those alternatives are the waste of mollusks and other aquatic organisms and the exoskeletons of crustaceans through the production of organic material (AL- Malikshah.2023.)

Chitosan has been used in several agricultural experiments and has shown clear superiority in vegetative indicators and yield, and these The material is Chitosan polymer (Sathiyabama *et al*, 2014).which is one of the substances found in the cell walls of the exoskeletons of arthropods, fungi, crustaceans, crabs, shrimp, and fish scales. Nano technology was used to produce this compound in its Nano form, which works to increase efficiency and effectiveness while increasing the surface area of the material so that it is added in a slow and easy release method, which provides a continuous supply of nutrients (Vurukonda *et al* , 2018). it has been widely used in the fields of biomedicine, biotechnology and agricultural engineering (Hidangmayum *et al* , 2019). The polymer also works to resist dryness through control It helps close stomata and reduce the water level in the leaves (Naeem *et al*, 2020) This substance helped induce resistance to diseases and stimulate the plant to produce effective, useful enzymes. For plants (Chun and Chandrasekaran, 2019),. It is also used to combat rot on fruit during canning and exporting from by covering the fruits with a thin layer of it (pan *et al* 2022). It is a biopolymer that can be obtained from Chitin (Abdulrasool and Al-Malikshah, 2022) Chitosan is known as a straight-chain copolymer

consisting of D-glucosamine and N-acetyl -D-glucosamine linked by  $\beta$ -type bonds (Al-Temimi, 2020)

Therefore, the research aimed to: test the efficiency of the best combination of Chitosan Nano fertilizer with the ground biological addition of *Azotobacter* bacteria for the purpose of increasing the vegetative indicators and yield of Kohlrabi plants.

## MATERIALS AND METHODS

An experiment was conducted during the fall semester of 2023 in one of the fields of the Plant Production Department in the desert region (the Tomato Development Project) of the Agriculture Directorate of the Holy Karbala Governorate. The seeds of the (hybrid) variety were planted on September 20, 2023, and care continued until seedlings with four true leaves were obtained. Kohlrabi seedlings were transferred to the field on October 25, 2023. The purpose of the research is to study the effect of two factors. The first factor is contamination of the roots of the plant seedlings with three levels of the *Azotobacter* biological additive (5 and 10) grams. A plant added to the no-addition treatment (Ahmad, 2024). The second factor is spraying the plant's foliage after 14 days of transplantation with Nano-Chitosan at a concentration of (0, 1, and 2) grams.  $L^{-1}$ . After two weeks of transplanting, two sprays every 14 days until complete wetness, using a portable sprayer with a capacity of 16 liters, drops of Sodium Tripolyphosphate were added as a dispersing agent to reduce the phenomenon of surface tension. The experiment followed a Randomized Complete Block Design (RCBD) with two factors and three different replications. with three replications and two factors (AL-Rawi and Khalf Allah, 2000). The number of transactions reached 27 experimental units. The means were compared according to Duncan's multiple range test at the probability level of 0.05. The seedlings were planted on terraces, each terrace width 75 cm and the distance between each terrace was 50 cm to isolate the treatments. The planting distance was 60 cm between one seedling and another and on both sides of the terrace to adopt dense planting. The length of each experimental unit was 3 m. The irrigation process was carried out using the drip irrigation method with 12 plants in the experimental unit. The irrigation process was carried out using the drip irrigation method. Five samples of field soil were taken from different places before planting, at different depths between 10-30 cm, then mixed well and a sample was taken to one of the private scientific laboratories for analyses. The results were as shown in Table (1).

**Table (1) Some physical and chemical characteristics of field soil**

Type of analysis	Unit of measurement	measurement
pH	—	7.8
EC	$ds^{-1}.m$	3.41
N	$mg.kg^{-1}$	17
P	$mg.kg^{-1}$	4.3
K	$mg.kg^{-1}$	59.9
$Na^{+}$	$mg.l^{-1}$	298.8
$SO_3$	$meq.l^{-1}$	573.7
$Cl^{-}$	$mg.l^{-1}$	339
$Ca^{++}$	$mg.l^{-1}$	229.42

The research included measuring the following indicators

### 1- The height of the plant (cm)

The lengths of the plants were measured from the soil surface to the end of the growths with a metric ruler and the average was taken.

### 2- Number of leaves per sheet. Plant<sup>-1</sup>

The number of leaves per plant was calculated by averaging five plants.

### 3- Total Chlorophyll content of leaves (mg 100 g<sup>-1</sup>)

Five plants were randomly selected. The leaves were taken from them and after washing them well from the dust (Al-Sahaf, 1989) . Using with acetone mash, a clear, white extract was obtained. Using a Spectrophotometer, optical absorption was measured at two wavelengths: 645 and 663, and the amount of chlorophyll was calculated according to the following equation.

$$[20.2 \times D(645)] + [8.02 \times D(663)] \times (v / w \times 1000) \times 100 = \text{Total chlorophyll}$$

### 4- The Weight of the knobs g plant<sup>-1</sup>

The knobs of five plants in each experimental unit were weighed and the average was taken

### 5- Total marketable yield

Small, infected and deformed knobs were excluded from the plant yield, and the remaining knobs were then calculated as a marketable yield according to the following:

$$\text{Total marketable yield (tons.h}^{-1}\text{)} = \frac{\text{Experimental unit yield} \times 10,000 \text{ m}^2}{\text{Experimental unit area}}$$

### 6- Percentage of nitrogen content in leaves: %

Total nitrogen (N) was determined by distillation using a Microkaldal.( Jackson,.1958)

The volume of acid consumed by elution x the standard of acid x 14 x the volume of dilution

$$N\% = \frac{\text{Volume of sample taken upon distillation} \times \text{weight of digested sample} \times 1000}{\text{Volume of acid consumed by elution} \times \text{the standard of acid} \times 14 \times \text{the volume of dilution}} \times 100$$

### 7- Protein of knobs %

The percentage of protein in the knobs was calculated on the basis of dry weight according to the following equation:

$$\text{Protein percentage based on dry weight} = \text{percentage of nitrogen in knobs} \times 6.25$$

Then it was calculated based on the wet weight as in the following equation:

$$\% \text{ of protein based on wet weight} = \frac{\% \text{ of protein based on dry weight} \times \% \text{ of dry matter in knobs}}{100} \quad (\text{A.O.A.C, 1980})$$

## RESULTS AND DISCUSSION

Table (2) shows the superiority of the treatment of inoculating the roots of the Kohlrabi seedlings with the *Azotobacter* at a level of 10 g in the plant height trait, as it gave the highest height of 47.33 cm. As for the effect of the single factor, spraying Nano chitosan at a concentration of 2 g L<sup>-1</sup>, it showed a significant superiority in the rate of plant height, reaching 54.56 cm compared to the comparison treatment. While the interaction between inoculating the roots of the Kohlrabi seedlings with the biological vaccine at a level of 5 g

and spraying nano chitosan at a concentration of  $2 \text{ g L}^{-1}$  in the plant height trait, as it gave the highest height of 60 cm compared to the untreated plants, which gave 34.33 cm .

Table (2) shows the superiority of the treatment of inoculating the roots of the Kohlrabi plant seedlings with the biological vaccine at a level of 5 g in the number of leaves in the plant, as it gave the largest number 17 leaves . As for the effect of the single factor, spraying Nano chitosan at a concentration of  $2 \text{ g L}^{-1}$ , it showed a significant superiority in the average number of plant leaves, reaching 19.56. compared to the comparison treatment. While the interaction between inoculating the roots of the Kohlrabi plant seedlings with the biological vaccine with spraying Nano chitosan in the number of leaves trait, as it gave the largest number of 21.67 . compared to the untreated plants, which gave the least number of plant leaves 9.33.

Table (2) shows the superiority of the treatment of inoculating the roots of the Kohlrabi plant seedlings with the bio- vaccine at a level of 10 g in the characteristic of the leaf content of chlorophyll as it gave the highest content 67.68. mg.  $100 \text{ g}^{-1}$  As for the effect of the single factor, spraying Nano-chitosan at a concentration of  $2 \text{ g L}^{-1}$ , it showed a significant superiority in the leaf content of chlorophyll mg.  $100 \text{ g}^{-1}$  74.16 compared to the comparison treatment. While the interaction between inoculating the roots of the Kohlrabi plant seedlings with the bio-inoculate at a level of 5 g and spraying Nano-chitosan at a concentration of  $2 \text{ g L}^{-1}$  in the characteristic of the leaf content of chlorophyll as it reached 77.59 mg.  $100 \text{ g}^{-1}$  compared to the untreated plants, which gave the lowest values 51.84 .

Table (3) shows the superiority of the treatment of inoculating the roots of the Kohlrabi plant seedlings with the biological vaccine at a level of 10 g In terms of plant weight, as it gave the largest corm weight 197.8 g As for the effect of the single factor, spraying Nano-chitosan at a concentration of  $2 \text{ g L}^{-1}$ , it showed a significant superiority in the corm weight reaching 251.3 g. compared to the comparison treatment. While the interaction between inoculating the roots of the Kohlrabi plant seedlings with the biological vaccine at a level of 5 g and spraying Nano-chitosan at a concentration of  $2 \text{ g L}^{-1}$  in the corm weight trait, as it reached 278.9 g . compared to the untreated plants, which gave the lowest values 78.4 g.

Table (3) shows the superiority of the treatment of inoculating the roots of the Kohlrabi plant with the biological vaccine at a level of 10 g in the total yield characteristic of , as it gave the highest yield of  $10.54 \text{ ton. ha}^{-1}$ . As for the effect of the single factor, spraying Nano-chitosan at a concentration of  $2 \text{ g L}^{-1}$ , it showed a significant superiority in the total yield of , as it gave the highest yield of  $13.40 \text{ ton. ha}^{-1}$  compared to the comparison treatment. While the interaction between inoculating the roots of the Kohlrabi plant with the biological vaccine at a level of 5 g and spraying Nano-chitosan at a concentration of  $2 \text{ g L}^{-1}$  in the total yield characteristic of as the yield reached  $14.87 \text{ ton. ha}^{-1}$  compared to the untreated plants, which gave the lowest values  $4.18 \text{ ton. ha}^{-1}$  .

Table (3) shows the superiority of the treatment of inoculating the roots of the Kohlrabi plant with the bio-vaccine at a level of 5 g in the characteristic of the nitrogen content of the leaves, as it gave the highest content of 2.74 % . As for the effect of the single factor, spraying Nano-chitosan at a concentration of  $2 \text{ g L}^{-1}$ , it showed a significant superiority in the nitrogen content of the leaves, as it gave the highest content of 3.10 % . compared to the comparison treatment. While the interaction between inoculating the roots of

the Kohlrabi plant with the bio-inoculate at a level of 5 g and spraying Nano-chitosan at a concentration of 2 g L<sup>-1</sup> in the characteristic of the nitrogen content of the leaves, as it reached 3.43 %. compared to the untreated plants, which gave lowest values 1.92% .

Table (3) shows the superiority of the treatment of inoculating the roots of the Kohlrabi plant with the biological vaccine at a level of 5 g in the characteristic of the protein content of the knobs %, as it gave the highest content of 17.14%. As for the effect of the single factor, spraying Nano chitosan at a concentration of 2 g L<sup>-1</sup>, it showed a significant superiority in the protein content of the knobs as it gave the highest content of 19.42% . compared to the comparison treatment. While the interaction between inoculating the roots of the Kohlrabi plant with the biological vaccine at a level of 5 g and spraying Nano chitosan at a concentration of 2 g L<sup>-1</sup> in the characteristic of the protein content of the knobs 21.48 %, as it reached compared to the comparison treatment, which gave 12.04 %.

**Table (2) The effectiveness of spraying with Nano-Chitosan and inoculating the roots of Kohlrabi seedlings with *Azotobacter* on Growth traits of plant and chlorophyll**

Treatment		height of plant cm <sup>-1</sup>	Number of leaves Leaves. Plant <sup>-1</sup>		Chlorophyll content of leaves (mg 100 g <sup>-1</sup> )	
Average of <i>Azotobacter</i> g plant <sup>-1</sup>	0	41.67 b	14.67	b	61.51	c
	5	47.11 a	17.00	a	66.33	b
	10	47.33 a	16.44	a	67.68	a
L.S.D		1.407	0.813		0.803	
Average of Chitosan g L <sup>-1</sup>	0	37.11 c	11.56	c	55.37	c
	1	44.44 b	17.00	b	65.99	b
	2	54.56 a	19.56	a	74.16	a
L.S.D		1.407	0.813		0.803	
<b>Azotobacter</b> <b>g plant<sup>-1</sup></b> <b>X</b> <b>Chitosan</b> <b>g L<sup>-1</sup></b>	0	34.33 g	9.33	f	51.84	h
	5	36.67 g	11.67	e	54.89	g
	10	40.33 f	13.67	d	59.38	f
	0	42.67 ef	15.67	c	61.85	e
	5	44.67 de	17.67	b	66.50	d
	10	46.00 cd	17.67	b	69.61	c
	0	48.00 c	19.00	b	70.84	c
	5	60.00 a	21.67	a	77.59	a
	10	55.67 b	18.00	b	74.06	b
L.S.D		2.436	1.408		1.391	

Means within a column, row and their interactions followed with the same letters are not significantly different from each other according to Duncan multiple ranges test at significant level of 5%.

**Table (3) Effectiveness of spraying with Nano-Chitosan and inoculating the roots of Kohlrabi seedlings with *Azotobacter* on quality traits and plant yield**

Treatment		The Weight of the knobs cm plant <sup>-1</sup>		The total yield (ton. h <sup>-1</sup> )		Percentage of nitrogen content in leaves: %		Protein content of knobs %		
Average of Azotobacter g plant <sup>-1</sup>	0	156.9	c	8.36	c	2.45	c	15.33	c	
	5	191.5	b	10.21	b	2.74	a	17.14	a	
	10	197.8	a	10.54	a	2.64	b	16.54	b	
L.S.D		1.161		0.064		0.092		0.577		
Average of Chitosan g L <sup>-1</sup>	0	107.7	c	5.74	c	2.03	c	12.72	c	
	1	187.1	b	9.97	b	2.70	b	16.88	b	
	2	251.3	a	13.40	a	3.10	a	19.42	a	
L.S.D		1.161		0.064		0.092		0.577		
Azotobacter g plant <sup>-1</sup> X Chitosan g L <sup>-1</sup>	0	0	78.4	i	4.18	h	1.92	f	12.04	f
		5	109.2	h	5.82	g	1.98	f	12.37	f
		1	135.6	g	7.23	f	2.20	e	13.75	e
		0	157.6	f	8.40	e	2.52	d	15.77	d
	1	0	186.3	e	9.93	d	2.81	bc	17.56	bc
		5	217.4	d	11.59	c	2.76	c	17.29	c
		1	234.8	c	12.52	b	2.91	bc	18.19	bc
		0	278.9	a	14.87	a	3.43	a	21.48	a
	2	5	240.3	b	12.81	b	2.97	b	18.58	b
		1								
		0								
		0								
L.S.D		2.011		0.110		0.160		0.999		

Means within a column, row and their interactions followed with the same letters are not significantly different from each other according to Duncan multiple ranges test at significant level of 5%.

We note from the tables (2) that spraying with the Nano-polymer chitosan significantly improves the characteristics (Tall of plant , number of leaves, chlorophyll content of the leaves). This may be attributed to the biological and Nano-components of the polymer, and this may contribute to the integration of cell division and recovery. Photocells reflect positive signals (Al-Khafaji , 2023) . The results showed a clear effectiveness of adding the bio fertilizer *Azotobacter*, which is attributed to the effectiveness of these organisms in the soil and the provision of a suitable environment in the area around the roots, in addition to the abundance of nutrients as a result of their metabolic processes. This has encouraged an increase in the efficiency of the roots and has been reflected in the growth of the root system and the size of the corm, and this is consistent with (Ahmad, 2024).

Tables 3 show a clear increase in the nitrogen content of the leaves and protein content of knobs. This may be due to the effectiveness of the biopolymer and its Nano property, as the polymer penetrates the cellular membranes easily and slowly. It has contributed to an increase in the synthesis of the chlorophyll molecule (Elshamy,2019), the basis of which is the porphyrin ring, thus increasing the efficiency of metabolism. Photosynthesis in plants, and the effectiveness and activity of microorganisms may have helped provide phosphorus to the roots, and this reflected positively on growth and yield (Fanaei et al., 2015 ,Caddell,



2019). The interaction between the two factors may have contributed to pushing the plant to grow and increasing yield because of their effect on the physiological aspect of the plant through the accumulation of organic matter and the availability of nutrients in the soil solution and the plant, and thus affects the biomass of the plant, and it reflects positively on the rest of the growth characteristics and yield. (Ahmad *et al.*, 2024).

Treatment with polymer nanoparticles has contributed to an increase in most vegetative growth indicators (plant height, number of leaves, leaf area, corm diameter). The reason for this may be attributed to the role of the Nano polymer because it contains the main and secondary elements, being an organic chitin us compound that contributes to the plant's vital activities and resistance to conditions. Environmental conditions, and may have stimulated the production of plant hormones such as cytokinin, gibberellin, and auxin (Mady, 2014). These compounds cause the plant to divide and elongate cells, thus leading to an increase in growth, and this in turn reflects positively on increased branching and the size of vegetative growth .

The Nano state of the compound is one of the factors of slow and continuous release, and this may have contributed to the longer-term supply of micronutrients, as nanoparticles are involved in the formation of proteins and amino acids and in the porphyrin ring that forms the chlorophyll molecule, which is one of the most important factors for plant growth and health, (Singh and Singh. 2008 ).

The use of chitosan polymer in its Nano form may contribute to improving the availability of phosphorus to the plant, which improves the condition Nutritional status of the plant and prevents The loss of nutrients in the soil also contributes to an increase in the activity of microorganisms in it.( Al-Malikshah, 2023).

## CONCLUSIONS

The sprayed with Nano-chitosan and inoculating the roots of the Kohlrabi plant with the biological vaccine (*Azotobacter*) with their interaction showed clear superiority in growth indicators, effectiveness in plant growth and improving soil parameters as a result of being organic materials first and being in the Nano state second.

## CONFLICT OF INTEREST

The authors declare no conflicts of interest associated with this manuscript.

## ACKNOWLEDGMENTS

The authors gratefully acknowledge the staff of the [Department of sciences, open educational College and to College of Education for Pure Sciences. Ibn Al-Haitham, University of Baghdad their technical and general support.

## REFERENCES

- A.O.A.C (1980) Official Methods of Analysis. 13<sup>th</sup> Ed. Association of Official Analytical Chemists. Washington. USA. D.C.
- Abdulrasool, I.J. and Z.R.J. Al-Malikshah (2022) Effect of Adding Fulzyme plus and Spraying with Green Tea Extract on Vegetative Growth and Yield of Pepper Cv.

- California Wonder. Basrah Journal of Agricultural Sciences.35(2) :302-312. <https://doi.org/10.37077/25200860>.
- Ahmed, G, A(2016) Efficiency of some antioxidants and bio agents in controlling Rhizoctonia Damping –off SnaBean. Middle East Journal of Applied Sciences 16:2.
- Ahmad.a RAH (2024)The Effect of Spraying with Plant Extracts on Some Growth Characteristics and Active Ingredients of Basil Plant. IOP Conf. Series: Earth and Environmental Science 1371 . doi:10.1088/1755-1315/1371/5/052003.
- Ahmad.b , R,A (2024)Effect of Mycorrhizal Inoculation and Fertilization with Plant Residues on the Growth of Chard Plant. Ibn AL- Haitham Journal For Pure and Applied Science 37(1):54-65. [doi.org/10.30526/37.1.3168](https://doi.org/10.30526/37.1.3168)
- Ahmad , R.A.H S.D.A.A. Twaij ,S.D.A.A, and Ibrahim ,A.M (2024) Phytohormones , Bio- And Mineral Fertilizers Effects on the Growth and secondary Compounds Of Chamomile (*Matricaria Chamomilla* L.) . SABRAO Journal of Breeding and Genetics. 56 (3) 1271-1281, 2024 doi.org/10.54910/sabrao2024.56.3.34.
- Al-Khafaji, A. M. H. H., and K. D. H. Al- Jubouri. (2023) Upgrading growth, yield, and folate levels of lettuce via salicylic acid and spirulina , vermicompost aqueous extracts,54(1):235-241 . <https://doi.org/10.36103/ijas.v54i1.1696>.
- Al-Bayati .A.S., H.A.M. Turk, A.K.H. AL-Tufaili , M.A. Aboohanah, R.K. Mohan, and H.M. Qader (2023) Characterization of green onion with NPK fertilization and Foliar Application of Hornwort Extract . SABRAO Journal of Breeding and Genetics .DOI: 10.54910/sabrao2023.55.6.25.
- Al- Sharia , Z.A.S (2022) Effect of spraying with marine Seaweed and Azotobacter treatment on growth yield and Content of Capsaicin compound for three cultivars of Chili pepper. A Thesis. Faculty of Agriculture University of Kufa. Ministry of Higher Education and Scientific Research. Iraq.
- Ali, A.H.A.A, Khalid A, M(2023) Effect of Bio health Bio stimulant and Dry Yeast Suspension on some Chemical Characteristics of Leaves and the Total Yield of Two Varieties of Kohlrabi Brassica oleracea var Gongylodes. IOP Conf. Series: Earth and Environmental Science 1158 (2023) doi:10.1088/1755-1315/1158/4/042060.
- Al-Juthery, H.W.A. and E.H.O. Al-Maamouri. (2020) Effect of Urea and Nano-Nitrogen Fertigation and Foliar Application of Nano-Boron and Molybdenum on some Growth and Yield Parameters of Potato. AlQadisiyah Journal for Agriculture Sciences.10(1):253-263. <http://qu.edu.iq/jouagr/index.php/QJAS/index>
- Al-Malikshah.Z.R.J. (2023) Effect of Spraying with Chitosan Nano Particles, Nettle leaf extract and Green tea extract on the Growth and Yield of Potato and Physical and Chemical Characteristics of Starch. PHD. Dissertation. Ministry of Higher Education. University of Baghdad. Iraq.
- Al-Mosawy.H.A.A(2023) Response of growth yield and quality indicators of two types of kale (*Brassica oleracea* var gongylodes) to the addition of the bio fertilizer (Bio health) and spraying bread yeast suspension. A Thesis. University of Karbala. Ministry of Higher Education and Scientific Research. Iraq.
- AL-Rawi, K.M. and A.M. Khalf Allah (2000) Design and Analysis of Agricultural Experiments. College of Agricultural . Univ. Mosel., Iraq.

- Al-Sahaf, Fadel Hussein (1989) Applied Plant Nutrition. Baghdad University, Ministry of Higher Education and Scientific Research.
- Al- Timimi, Q.A.A(2020) Test of the efficacy of Nano Chitosan in the resistance of early blight on Tomato caused by the Fungus *Alternaria solani*. A Thesis. College of Basic educational Al- Mustansiriyah University. Ministry of Higher Education and Scientific Research. Iraq.
- Caddell.D.F, Siwen .D, Devin. C(2019) Seed Endophytes Biology and Biotechnology Cha 1 . April 2019.pp 273–311
- Chun. S.C, Chandrasekaran, M (2019) Chitosan and Chitosan nanoparticles induced expression of pathogenesis related protein genes enhances biotic stress tolerance in tomato. International journal Biological Macromol .125: 948-954.
- Elshamy, M.T.; S.M. Elkhallal; Sh.M. Hussein and K.Y. Farroh (2019) Application of Nano-chitosan NPK fertilizer on growth and productivity of potato plant. Journal of Scientific Research in Science. 36(1):424-441.
- Fanaei , H.R, Hanieh. K, Eisa .P (2015) Effect of seed priming on grain and oil yield of Safflower under irrigation deficit conditions. Iranian Journal of Seed Science and Research, 2015, Vol. 2, No. 2, P49- 59 .
- Hidangmayum, A, Dwivedi, P, Katitar, D, and Hemantaranjan, A (2019) Application of chitosan on plant responses with special references to abiotic stress. Physiology and molecular biology of plants .25(2): 313-326.
- Jackson, M.L .1958. Soil Chemical Analysis. Prenticahall Inc Englewood Cliffs. N.J.183-204.
- Mady , M. A(2014) Inducing cold tolerability in squash (*Cucurbita pepo* L.) plant by using salicylic acid and chelated calcium application . Inter. J. of Agric. Sci. and Research. 4 (4) : 9-24
- Mohammed.M. S, Syeda.J.T.M, Wasan, K.M.and Wasan.E.K(2017)An overview of chitosan nanoparticles and its application in nano-parenteral drug delivery. pharmaceutics .9: 53.
- Naeem U, A, Basit, I, Ahmad, I, Ullah, S, T, Shah, H, I. Mohamed and S, Javed (2020) Mitigation the adverse effect of salinity stress on the performance of the tomato crop by exogenous application of chitosan. Bulletin of the National Research Centre.Vol 44, 181.
- Pan , Q.; C. Zhou; Z. Yang; Z. He; C. Wang; Y. Liu; S. Song; H. Gu; K. Hong; L.Yu; Y.Qu and P.Li(2022) Preparation and Characterization of Chitosan derivatives modified with quaternary ammonium salt and quaternary phosphate salt and its effect on tropical fruit preservation. Food Chemistry. 387:132878.
- Sabalpara, N, A, Lalit, M, (2016) Effective Utilization of Bio fertilizers in Horticultural Crops. Commercial Horticulture, New Delhi, India. pp 185-192.
- Sathiyabama, M. Akile. G, and Charel, R.E(2014)Chitocan induced defence responses in tomato plants against early blight disease caused by *Alternaria solani* (Ellis and Martin) Sorauer.Achives of phytopathology and plant protection. 47 (16): 1963-1973.
- Singh, A. and P. K. Singh .2008. Salicylic acid induced biochemical changes in cucumber cotyledons. Indian J. of Agricultural Bio ch. 21(1and2), 35 – 38.

- Taiz, L. and E. Zeiger (2014) Plant Physiology. 5rd.ed Sinauer Associates, Inc., Publishers. Sunderland, Massachusetts.
- Turan M, Ataoglu N, and Sahin F. 2006. Evaluation of the capacity of phosphate solubilizing bacteria and fungi on different forms of phosphorus in liquid culture. J. Sustain. Agr., 28(3):99–109.
- Turk, H.A, A(2023) Response of Two Potato Cultivars to Spraying with BiozymeTF and *Azosprillum* on some Growth and Yield Indicators. IOP Conf. Series: Earth and Environmental Science 1158 (2023) 042033. doi:10.1088/1755-1315/1158/4/042033.
- Turk, H.A, Mansour. A.A, Mahmood. T.M, (2020) The Effect of Adding Animal Manure and Bio Stimulant Amalgerol on the Growth Parameters of two Verities (Local, Spanish) of the Broad Bean Plant (*Vicia Faba* L.). Int. J. Agricultural. Stat. Sci. pp. 1365-1370.
- Vurukonda, S.S.K.P ; Giovanardi , D; and Stefani ,E (2018) Plant Growth Promoting and bio control activity of *Streptomyces* spp. And endophytes . international journal of molecular sciences .19(4): pp 952.